

1998

Decline of Iowa Populations of the Regal Fritillary (*Speyeria idalia*) Drury

Diane M. Debinski

Iowa State University, debinski@iastate.edu

Liesl Kelly

Iowa State University

Let us know how access to this document benefits you

Copyright © Copyright 1998 by the Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/jias>



Part of the [Anthropology Commons](#), [Life Sciences Commons](#), [Physical Sciences and Mathematics Commons](#), and the [Science and Mathematics Education Commons](#)

Recommended Citation

Debinski, Diane M. and Kelly, Liesl (1998) "Decline of Iowa Populations of the Regal Fritillary (*Speyeria idalia*) Drury," *Journal of the Iowa Academy of Science: JIAS*, 105(1), 16-22.

Available at: <https://scholarworks.uni.edu/jias/vol105/iss1/4>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Journal of the Iowa Academy of Science: JIAS by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Decline of Iowa Populations of the Regal Fritillary (*Speyeria idalia*) Drury

DIANE M. DEBINSKI and LIESL KELLY

Dept. of Animal Ecology, Iowa State University, Ames, Iowa 50011
email: debinski@iastate.edu

The Regal Fritillary butterfly, *Speyeria idalia* Drury (Lepidoptera: Nymphalidae, Argynninae), is a prairie endemic species which has exhibited a dramatic decline in Iowa during the last few decades. Tallgrass prairie is the primary habitat of *S. idalia* and the butterfly's presence is correlated with the presence of violets (*Violaceae*). Due to the extensive habitat fragmentation of Iowa prairies, both the butterfly and its host plant populations are limited primarily to unplowed, relatively pristine prairies. Thus, *S. idalia* is an excellent case study of a prairie endemic species. Here, we report on the results of a two-year survey of the insect and its host plants in Iowa. During 1995, we conducted extensive surveys of 52 prairies in southern Iowa and found *S. idalia* in 11 of those prairies. During 1996, we conducted more intensive surveys comparing eight Iowa prairies to prairies in Kansas, South Dakota, and North Dakota to examine the hypothesis that larval host plant limitation may be causing the decline of *S. idalia*. Our data show that Iowa prairies have lower hostplant availability and also have lower butterfly weights. We discuss the future prospects for the species and suggest various management scenarios that might aid in preserving the butterfly.

INDEX DESCRIPTORS: *Speyeria idalia*, butterfly, *Violaceae*, prairie, habitat fragmentation, extirpation

HISTORICAL PERSPECTIVE

Tallgrass prairies in the U.S. exemplify one of the most highly fragmented natural areas, with 1–12% of their original area remaining (Sampson and Knopf 1994). In Iowa, less than 0.02% of these habitats remain intact (Smith 1981). One of the most visible components of prairie biodiversity is the butterfly community, and the Regal Fritillary (*Speyeria idalia* Drury Lepidoptera: Nymphalidae, Argynninae) might well be considered a symbol of prairie endemic insects. Tallgrass prairie is the primary habitat of *S. idalia* (Hammond and McCorkle 1983, Opler and Krizek 1984, Schull 1987) and the butterfly's presence is correlated with the presence of violets (*Violaceae*). *S. idalia*'s larval host plants include *Viola pedata* (Bird's-foot Violet), *V. pedatifida* (Blue Prairie Violet), *V. papilionacea* (Common Blue Violet), *V. lanceolata* (Lance-leaved Violet) and *V. nuttallii* (Nuttall's Violet) (Schull 1987, Opler and Krizek 1984). Regions in Iowa where *S. idalia* is most abundant contain largely *V. pedatifida*. With the disappearance of prairie habitat, widespread populations of *S. idalia* also have declined in numbers and distribution. This trend is especially pronounced in Iowa due to the intensity of agriculture. Larger populations of *S. idalia* can be found in Great Plains states that have larger prairies remaining.

Historic records of the species in Iowa date back to Scudder's surveys in 1869. Records from 1869 to the present reveal that the species was known to occur in at least 51 of the 98 Iowa counties (Schlicht pers. com.). It may have been even more common. Population estimates of this insect in Iowa and across the nation have declined sharply over the last 50 years, primarily due to plowing of the remaining prairies. *S. idalia* was listed as a Category II species under the Endangered Species Act until 1996, when this category of protection was deleted by the U.S. federal government (USFWS 1996). Category II species were candidates for listing, but there was not sufficient knowledge regarding their status to warrant proposing them for listing as endangered or threatened. *S. idalia* currently is

listed as a species of special concern in Iowa and in other prairie states (J. Fleckenstein, pers. com.).

METHODS

In 1995, we conducted an extensive survey of *S. idalia* and *V. pedatifida* populations in southern Iowa (Fig. 1). Using knowledge of the species' historic distribution and information regarding potential habitats remaining in southern Iowa, we surveyed 52 prairies. The butterfly was surveyed using standard mark-recapture techniques (Pollard 1977) and Lincoln-Peterson calculations (Davis and Winstead 1980). The violets were surveyed for presence/absence in all prairies, and *V. pedatifida* density was examined relative to prairie management history on 29 of the 52 prairies. In 1996, we conducted more intensive surveys of *S. idalia* populations and *V. pedatifida* densities, focusing on eight of the Iowa prairies that had large *S. idalia* populations and comparing these prairies to prairies in Kansas, Nebraska, and North and South Dakota.

In 1995, our *V. pedatifida* searches lasted for a minimum of 30 min, unless the habitat was obviously poor quality [e.g., solid *Panicum virgatum* (switchgrass), *Bromus* spp. (brome), etc.]. The densest patches of *V. pedatifida* were located at each prairie. A 10 x 10 m plot was centered in each of these areas, and three plots were surveyed in most prairies (except where prairie size limited sampling). *V. pedatifida* density was estimated by counting the number of plants (stems with a single root stalk) in every other square meter in each of 10 transects that made up the plot, and mean *V. pedatifida* density was calculated across all plots. To understand the effects of management on *V. pedatifida*, each prairie was ranked relative to the intensity at which it experienced a given land use (ranging from 0 to 3). Land use categories included mowing, burning, and grazing. Each prairie was also ranked according to its moisture level (dry, mesic, or wet).

In 1996, we examined the issue of host plant limitation more intensively by focusing our surveys on fewer, more high quality prairies (Fig. 2). We compared Iowa data on butterflies and host plants




Fig. 1. Iowa prairies surveyed for *S. idalia* in 1995. Filled circles are sites where the species was found and open circles are sites where the species was not found. Site codes are as follows: 100: Adair Wildlife Area, 101: Honey Creek State Park, 102: Boone Railroad Prairie, 103: Troy Township Cemetery, 104: Flaherty Prairie, 105: Page Private Prairie, 106: Little River Recreation Area, 107: Nine Eagles State Park, 108: Sand Creek Wildlife Area, 109: Slip Bluff Park, 110: Howe Prairie, 111: Sheeder Prairie, 112: Gleason-Hubel Wildlife Area, 113: Pioneer State Forest, 114: Murray Hill Overlook, 115: Kellogg Wildlife Area, 116: Kish-ke-kosh State Preserve, 117: Reichelt Unit, 118: Potato Creek, 119: Morris Prairie, 120: Linger Longer Prairie, 121: Stephens State Forest, 122: Lucas Railroad Prairie, 123: Johnson Prairie, 124: Chris Cove Park, 125: Peebler Prairie, 126: Hull Wildlife Area, 127: Hawthorn Wildlife Area, 128: Wearin Prairie, 129: Loess Hills Wildlife Area, Section 21, 130: Loess Hills Wildlife Area, Section 9, 131: Loess Hills Wildlife Area, Section 34, 132: Melrose Cemetery, 133: Pike Run Wildlife Area, 134: Red Cedar Wildlife Area, 135: Highway 70 Prairie, 136: Shield Prairie, 137: Kalsow Prairie, 138: Polk City Cemetery, 139: Moekley Prairie, 140: Saylorville's Red Feather Prairie, 141: Big Creek Prairie, 142: Wabash Trace Railroad Trail, 143: Malcom Railroad Prairie, 144: Ringgold Wildlife Area, 145: Doolittle State Preserve, 146: Raymond-Rolling Prairie, 147: McCallsburg Railroad Prairie, 148: Cumming Railroad Prairie, 149: Rolling Thunder Prairie, 150: Lake Darling, 151: Denison Prairie.




Fig. 2. Iowa prairies surveyed for *S. idalia* in 1996. The species was found in all sites and a more extensive survey of the butterfly and its host plant were conducted. Site codes are as follows: 111: Sheeder Prairie, 117: Reichelt Unit of Stephens State Forest, 129: Loess Hills Wildlife Area, Section 21, 137: Kalsow Prairie, 144: Ringgold Wildlife Area, 149: Rolling Thunder Prairie, 152: Cayler Prairie, 153: Anderson Prairie.

Table 1. *Speyeria idalia* population size estimates in 1995.

| PRAIRIE NAME | POPULATION ESTIMATE | METHOD |
|--|---------------------|------------------------------|
| Page Private Prairie | 2 | individuals observed |
| Sheeder Prairie | 50 | mark-recapture |
| Reichelt Unit of Stephens State Forest | 4 | individuals observed |
| Polk City Prairie | 1 | individuals observed |
| Moeckley Prairie | 220 | mark-recapture |
| Ringgold Wildlife Area | 7 | individuals observed |
| Doolittle Prairie | 2 | individuals observed |
| Rolling Thunder Prairie | 120 | mark-recapture |
| Kalsow Prairie | 500 | visual estimate ¹ |
| Loess Hills Wildlife Area sect. 9 | 160 | mark-recapture |
| Loess Hills Wildlife Area sect. 21 | 2 | individuals observed |

¹P. C. Hammond's estimate

with data from Kansas, South Dakota, and North Dakota prairies, where host plant abundance is greater. We surveyed five plots, rather than three plots for violets, and included *V. pedatifida* as well as other *Viola* species because *V. nuttallii* is much more common as a host plant in the Dakotas. In addition, we measured the size and weight of *S. idalia* butterflies in Iowa prairies and compared these values to violet density (for details, see Kelly 1997).

RESULTS AND DISCUSSION

In 1995, *S. idalia* was found in 11 out of the 52 prairies (Fig. 1), but only five prairies had a population over 50 individuals (Table 1). We found *S. idalia* only on sites where *V. pedatifida* were found, and population sizes of the insect seemed correlated with the violet biomass. *V. pedatifida* was found in 18 prairies. We were able to obtain information regarding management history on 29 of the 52 prairies, and this included 16 of the 18 prairies with *V. pedatifida* (Table 2). No *V. pedatifida* were found in any of the areas that had been plowed. These data support the assumption that plowing has a negative impact on *V. pedatifida*. Interestingly, violet densities were highest in moderately grazed and recently burned prairies, and the correlation between violet density and grazing intensity was significant (Spearman rank correlation = 0.43, $P = 0.02$). Grazing and burning probably release the plant from competition with many of the tallgrass species, offering it an opportunity to flourish. Moisture and violet density were negatively correlated (Spearman rank correlation = -0.43, $P = 0.06$), which is not surprising, given that the plants are usually found on dry hillsides. There was no significant correlation between mowing or burning and violet density.

V. pedatifida densities on Iowa prairies were not very high, averaging 1.85 plants/m² in 1995 (Table 2). In contrast, *V. papilionacea* can be found in densities of up to 20 stems/m² in Iowa forests (Kelly and Debinski, unpublished data). *Speyeria cybele*, a woodland butterfly species whose larvae feed on *V. papilionacea*, is found throughout Iowa. Given the fact that both *S. idalia* and *S. cybele* are large insects and consume a large biomass of violet leaves during the larval stage, a tenfold difference in host plant abundance could explain the relative rarity of *S. idalia* on Iowa prairies as compared to the abundance of *S. cybele*.

Our investigation of hostplant availability relative to *S. idalia* population size provided mixed results. In 1996, we found that North and South Dakota and Kansas had much larger violet populations (Table 3). However, there was not a clear relationship between violet population size and *S. idalia* population size. Iowa prairies showed lower *S. idalia* population sizes and violet densities than the Dakotas. However, the Kansas populations of *S. idalia* were actually lower

than the Iowa average, despite having a hostplant population that was six times larger.

Examining the hostplant limitation hypothesis from the perspective of insect weight provided support to our hypothesis. Iowa butterflies had statistically significant lower weights among both males and females, in contrast to the prairies we surveyed in Kansas, North Dakota, and South Dakota (Kelly 1997). Another striking result of these surveys was that we found adult sex ratios were almost 10 males to 1 female in Iowa prairies, whereas in other states the sex ratio was barely skewed. Because males emerge before females in this species (Opler and Krizek 1984), if males consume larval hostplant resources faster than females, females may lack sufficient resources to finish their larval stage. This may explain why the Iowa populations we observed had so few females. It may also be an important clue as to why Iowa *S. idalia* populations are declining. If prairie size limits the total violet biomass available, larval males may use up the resource in small prairies. Male populations of *S. idalia* were correlated (but not significantly) to area of habitat surveyed ($r^2 = 0.75$, $P = 0.17$, $df = 11$). Thus, there may be a prairie size threshold below which the populations are at a high risk of extirpation due to demographic problems precipitated by host plant limitation (e.g., Gilpin and Soule 1986).

FUTURE POSSIBILITIES AND POTENTIAL: CORRIDORS AND RESTORATION

Butterflies such as *S. idalia* are keenly affected by landscape patchiness and therefore provide a useful model system for examining the effects of prairie fragmentation in Iowa. Mechanisms resulting in such sensitivity include host plant specificity, physiological constraints, limited dispersal abilities of larvae, predator avoidance (Wood and Samways 1991, Litsinger et al. 1991, Thomas and Harrison 1992, Thomas et al. 1992, Rodrigues et al. 1993, Rodrigues et al. 1994), and even avoidance of crossing a prairie edge to move into a new habitat type (Ries and Debinski, unpublished data). These behavioral mechanisms lead to differential use and differing population sizes on prairie remnants. Subtle implications of these differences may even include effects on plant community dynamics such as altered herbivore loads, or changes in pollination (Hendrix, pers. comm.).

Although *S. idalia* is at less of a risk of extirpation across the entire Great Plains, our surveys of population sizes of the insect in Iowa during 1995 and 1996 suggest that it is at risk of extirpation in Iowa. Hammond and McCorkle (1983) attribute the decline of a number of *Speyeria* populations to the extent of environmental disturbances caused by humans. In Iowa prairies, the erratic distribution

Table 2. A synopsis of land use history and prairie characteristics of 29 Iowa prairies. Prairie violet density was calculated by averaging all plots from a prairie site. The number of plots taken from a site varied from 1 to 5 depending on the size of the prairie. Information on the soil moisture of the prairies was obtained from the Iowa Department of Natural Resources.

| SITE NAME | AVERAGE VIOLET DENSITY (#/m ²) | GRAZING | MOWING | BURNING | SOIL MOISTURE |
|--------------------------|---|----------------------------|-----------------------|----------------------------|------------------|
| Boone Railroad | 0 | none | none | accidental distant past | mesic |
| Little River Recreation | 0 | infrequent distant past | none | infrequent | mesic |
| 9 Eagles State Park | 0 | distant past | none | infrequent | dry |
| Howe Prairie | 1.9 | intensive | none | none | dry |
| Sheeder Prairie | 0.8 | none | annual until 1965 | once every three years | mesic |
| Gleason-Hubel | 0 | none | none | infrequent | dry |
| Pioneer State Forest | 0 | none | none | once every three years | dry |
| Murray Hill Overlook | 0 | infrequent distant past | none | infrequent | dry |
| Kellogg Wildlife | 0 | none | none | none | mesic |
| Kish-kekosh Prairie | 6.7 | lightly before 1976 | none | once every four years | dry |
| Reichelt Unit Prairie | 2.0 | moderate in past | none | once (1994) | dry |
| Potato Creek Prairie | 0 | moderate to intensive | moderate | once every 5(+) years | mesic |
| Morris Prairie | 0.1 | none | none | once every 3-4 years | mesic |
| Hull Wildlife | 0 | none | none | none | mesic |
| Hawthorne Wildlife | 4.5 | none | none | once every five years | mesic |
| Loess hills Wildlife - 9 | 2.9 | intensive | none | infrequent distant past | dry |
| Loess hills Wildlife-21 | 1.6 | moderate in past | none | once every five years | dry |
| Loess hills Wildlife-34 | 0.1 | none | none | infrequent distant past | dry |
| Polk City Cemetary | 0.1 | none | none | once every three years | dry |
| Moeckley Prairie | 0.7 | none | annual before 1987 | once | wet |
| Saylorville Prairie | 0 | none | none | once every five years | wet |
| Big Creek Prairie | 0 | none | moderate in past | once every three years | mesic |
| Wabash Trace Trail | 0 | none | none | accidental distant past | mesic |
| Ringgold Wildlife | 0.5 | moderate | none | once every five years | dry |
| Doolittle Prairie | 0.1 | moderate | moderate | once every four years | wet |
| Raymond/Rolling | 4.6 | light | none | once every five years | dry |
| McCallsburg Railroad | 0 | none | none | once every 3-5 years | mesic |
| Cummings Railroad | 1.1 | none | none | once every 2-3 years | mesic |
| Rolling Thunder | 1.9 | intensive distant past | none | once every 3-4 years | dry |

Note: Page Private Prairie and Kalsow Prairie are not included in the land use history survey, but they did have *V. pedatifida* present in 1995.

Table 3. *Speyeria idalia* population size estimates relative to violet densities in Iowa prairies as compared to North and South Dakota and Kansas Prairies in 1996.

| SURVEY LOCA- TIONS | | HEC- TARES SUR- VEYED | VIOLET POPU- LATION ESTIMATE | VIOLET DENSITY PLANTS/m ² | S. IDALIA 1996 POP. ESTI- MATE | INSECT DEN- SITY EST. (INSECTS/ HA) |
|-----------------------------------|-------------------------|--------------------------------|---------------------------------------|--|---|---|
| Iowa | Loess Hills, Section 21 | 20.2 | 67,500 | 3,342 | 175 | 8.7 |
| | Sheeder Prairie | 10.1 | 22,600 | 2,238 | 163 | 16.1 |
| | Stephens State Forest | 8.1 | 13,500 | 1,667 | 21 | 2.6 |
| | Ringgold Wildlife Area | 64.8 | 86,000 | 1,327 | 270 | 4.2 |
| | Cayler State Preserve | 64.8 | 57,500 | 887 | 319 | 4.9 |
| | Anderson State Preserve | 64.8 | 57,400 | 886 | 494 | 7.6 |
| | Kalsow State Preserve | 64.8 | 136,000 | 2,099 | 224 | 3.5 |
| | Rolling Thunder | 49.8 | 423,000 | 8,494 | 60 | 1.2 |
| Iowa means | n = 8 | 43.4 | 107,938 | 2,617 | 216 | 6.1 |
| Dakota means (North and South) | n = 6 | 64.8 | 1,086,500 | 16,767 | 431 | 6.7 |
| Kansas means | n = 2 | 35.9 | 622,150 | 12,469 | 98 | 14.2 |

as well as low abundance of *V. pedatifida* is probably responsible for the small *S. idalia* populations. The primary cause of the restricted distribution of *V. pedatifida* in Iowa prairies is plowing. Plowing has caused serious fragmentation of prairie habitats, leaving *S. idalia* populations highly isolated. Grazing could actually have a positive, albeit indirect, effect on *S. idalia* populations via the effects on *V. pedatifida*. However, the intensity of the grazing would definitely have to be moderate. We cannot speculate here about effects of fire on *S. idalia*, but this is a research area that warrants further attention.

In the context of prairie management, many of the sites we surveyed in Iowa also had extensive areas of invasive brome with few if any violets. *S. idalia* adults often patrol these areas and feed at the nectar sources present in them. Such areas with no violets may be detrimental to female fecundity [i.e., sink populations (Pulliam 1988)] if adult females spend significant time searching there in vain for areas with host plants where they could deposit eggs.

In Iowa, we face three challenges in conserving this species. First, how do we manage and preserve the areas where the insect is consistently found? Second, how do we reconnect fragmented prairie remnants, and third, how do we enhance potential habitat for the insect where it once was found and/or could be reintroduced? In order to preserve the current populations of *S. idalia* in Iowa prairies, disturbances such as plowing and spraying on native prairies should be proscribed. Plowing destroys the root mass of violets, and additionally, *V. pedatifida* is a species that does not propagate well by seed (Debinski and Kelly, unpublished data). Unnecessary collecting of insects (especially females) should also be discouraged, since we found only 31 female insects across eight of the largest populations in the state in 1996.

There may be the potential to re-connect some prairie habitats through the use of corridors. One example of a possible corridor for *S. idalia* adults may be roadside planting of native prairie forbs and grasses. These roadsides could serve as important nectar sources and potentially aid in linking isolated prairie remnants. However, roadside plantings will probably never serve as good larval habitat due to the difficulty of establishing host plant populations from seed. Restoration of large prairies, such as the project at Walnut Creek

National Wildlife Refuge, offers some hope for *S. idalia*, but in order to establish a viable population, a large amount of financial and human resources would need to be committed to restore tens of acres of the violet host plants. It is undeniably a difficult undertaking to transplant and physically improve prairie habitat, but we (the authors) have participated in this effort, and have seen some grasslands improve in quality.

Finally, we need to assess the extent of decline as well as the reasons why *S. idalia* populations are disappearing in our prairies. We should monitor larger areas to determine possible reasons for population fluctuations of this species and initiate a citizen-based monitoring effort. Some of the larger areas that should be monitored include Ringgold Wildlife Area, in Ringgold County, Steele Prairie, in Cherokee County, and Stephens State Forest (Reichelt Unit), in Jasper County. In Ringgold Wildlife Area, there are over 1,200 acres of wildflowers to support adult insects with nectar plants, but there is only a very small *V. pedatifida* population, since much of the area was once plowed or otherwise disturbed. For this reason, the Ringgold Wildlife Area could be a good candidate for restoring violet populations. In the Reichelt Unit of Stephens State Forest, the situation is the opposite: there is quite a large population of *V. pedatifida*, but few nectar sources to maintain adult insects at that site. It is puzzling to us that a large resource base of violets (hundreds of thousands of plants) would not result in populations of adult insects larger than the 50 or so that we estimated in 1995 and 1996. In Steele Prairie, a state preserve, previous state records indicate that hundreds of insects were found regularly in the 1980s, but during the last season we surveyed (1996), we found only a few insects despite extensive violet coverage. These interesting survey results continue to demand the attention of conservationists in Iowa. We hope to initiate a program in the near future to ask outdoor enthusiasts to report when and where they find *S. idalia*. This would aid in cultivating local appreciation of rare Iowa wildlife and it would provide a much more extensive database from which to evaluate the status of *S. idalia* in the next decade.

ACKNOWLEDGEMENTS

We would like to extend our appreciation to the funding sources that made this research possible: the Iowa Science Foundation, the Iowa State Preserves Board, the Iowa Department of Natural Resources, and Iowa State University. Laura Myers, Eric Gundlach, and Jim Kolbet assisted in the field research. John Fleckenstein was helpful in suggesting prairies to include in our survey. Tim Orwig and Dennis Schlicht provided comments on earlier versions of this paper. This is Journal Paper No. J-17430 of the Iowa Agriculture and Home Economics Experiment Station, Ames, Iowa. Project No. 3377.

LITERATURE CITED

- DAVIS, D. E. and L. W. RAY. 1980. Estimating number of wildlife populations. Pages 221–246 *In*: Wildlife Management Techniques Manual. Schemnitz, S.D. (ed.). The Wildlife Society, Washington, D.C.
- GILPIN, M. E. and M. E. SOULE. 1986. Minimum viable populations: processes of species extinction. Pages 19–34. *In*: Conservation Biology: the science of scarcity and diversity. Sinauer Associates Sunderland, Massachusetts.
- HAMMOND, P. C. & D. V. MCCORKLE. 1983. The decline and extinction of *Speyeria* populations resulting from human environmental disturbances (Nymphalidae: Argynnninae). *Journal of Research on the Lepidoptera* 22: 217–224.
- KELLY, L. 1997. Exploring larval food limitation as a probable cause of decline in Iowa populations of a butterfly, *Speyeria idalia* (Lepidoptera: Nymphalidae). M.S. Thesis. Iowa State University.
- LITSINGER, J. A., V. HASSE, A. T. BARRION, and H. SCHMUTTERER. 1991. Response of *Ostrinia furnacalis* (Guenée) (Lepidoptera: Pyralidae) to intercropping. *Environmental Entomology* 20:988–1004.
- OPLER, P. A. & G. O. KRIZEK. 1984. Butterflies East of the Great Plains, an Illustrated Natural History. John Hopkins University Press, Baltimore.
- POLLARD, E. 1977. A method for assessing changes in the abundance of butterflies. *Biological Conservation* 12:115–134.
- PULLIAM, R. 1988. Sources, sinks, and population regulation. *American Naturalist* 132:652–661.
- RODRIGUES, J. J. S., K. S. BROWN, Jr., and A. RUSZCZYK. 1993. Resources and conservation of neotropical butterflies in urban forest fragments. *Biological Conservation* 64:3–9.
- RODRIGUES, J., D. JORDANO and J. F. HAEGER. 1994. Spatial heterogeneity in a butterfly-host plant interaction. *Journal of Animal Ecology* 63:31–38.
- SAMPSON, F. and F. KNOPF. 1994. Prairie conservation in North America. *BioScience* 4:418–421.
- SCHULL, E. M. 1987. The Butterflies of Indiana. Indiana Academy of Science.
- SMITH, D. D. 1981. Iowa prairie—An endangered ecosystem. *Proceedings of the Iowa Academy of Science* 88:7–10.
- THOMAS, J. A. 1983. A quick method for estimating butterfly numbers during surveys. *Biological Conservation* 27:195–211.
- THOMAS, C. D. and S. HARRISON. 1992. Spatial dynamics of a patchily distributed butterfly species. *Journal of Animal Ecology* 61:437–446.
- THOMAS, C. D., J. A. THOMAS, and M. S. WARREN. 1992. Distributions of occupied and vacant butterfly habitats in a fragmented landscape. *Oecologia* 92:563–567.
- U.S. FISH AND WILDLIFE SERVICE, UNITED STATES DEPARTMENT OF THE INTERIOR. 1996. 50 CFR Part 17, Endangered and Threatened Wildlife and Plants; Review of Plant and Animal Taxa That Are Candidates for Listing as Endangered or Threatened. Federal Register, Vol. 61(40):7596.
- WOOD, P. A. and M. J. SAMWAYS. 1991. Landscape element pattern and continuity of butterfly flight paths in an ecologically landscaped botanic garden, Natal, South Africa. *Biological Conservation* 58:149–166.